



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for Go Clear Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

1: J Orthop Res. 1996 Jul;14(4):573-81.

Related Articles, Link

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Spatial distribution of CD44 and hyaluronan in the proximal tibia of the growing rat.**Noonan KJ, Stevens JW, Tammi R, Tammi M, Hernandez JA, Midura RJ.**

Department of Orthopaedic Surgery, University of Iowa, College of Medicine, Iowa City 52242, USA.

CD44 has been described as a cell surface hyaluronan receptor present on a variety of different cells, and it is generally assumed to be prevalent in most connective tissues that contain hyaluronan. A major aim of this study was to test that presumption by localizing CD44 and hyaluronan within several tissues of the proximal tibia of the growing rat. Comparison of these profiles would reveal whether CD44 and hyaluronan co-localize with high fidelity, as would be expected if CD44 were a major hyaluronan binding protein. Using in situ hybridization and immunohistochemistry, CD44 was identified on osteoclasts, chondroclasts, osteocytes, hematopoietic marrow cells, synovial cells, and connective tissue fibroblasts (ligaments, tendons, and fascia). Although the majority of osteocytes expressed CD44, reduced expression was observed for osteoblasts and osteoprogenitor cells. Additionally, CD44 was not detected on chondrocytes from epiphyseal and metaphyseal growth cartilages or in meniscal fibrocartilage. Using biotinylated G1 domain from aggrecan and link protein, hyaluronan was observed in the maturational and hypertrophic zones of all growth cartilages, the synovium and other fibroblastic connective tissues, regional areas of the periosteum and endosteum (around osteoblasts, osteoprogenitor cells, and osteoclasts), osteocyte lacunae, and surrounding blood vessels. In regions of co-localization for CD44 and hyaluronan, it seems that CD44 is a likely hyaluronan binding protein in several tissues of the proximal tibia. However, it does not appear to be the predominant hyaluronan binding protein in growing cartilages of the weanling rat.

PMID: 8764866 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text